

Version 2 SOLUTIONS

Name:

ID number:

Time: 1 hour

Math 204

8/04/06

Quiz II

Second Semester 05/06

Instructor: Mrs. Muna Jurdak

Section 1: Thurs. 11:00 a.m.

Section 2: Fri. 1:00 p.m.

Section 3: Thurs. 3:30 p.m.

Section 4: Thurs. 2:00 p.m.

Instructions:

1. Write your name and ID number clearly where indicated.
2. Circle your section number above, according to the time of the problem solving session in which you are enrolled.
3. **Solve the problems on this, the white question sheet.** Use the colored sheets for scratch work only. You may use the back of a white sheet to complete the solution of a problem.
4. Give numerical answers correct to 2 decimal places.
5. **Show your work** in all the problems.

1	2	3	4	5	6	7	8	Total
/12	/10	/10	/15	/8	/17	/10	/18	/100

(12%) 1. (a) Find $\frac{dy}{dx}$ for $y = (e^{x^{3/2}}) \ln(x^3 + 1) = u \cdot v \rightarrow \frac{dy}{dx} = u'v + uv'$

$$\frac{dy}{dx} = \frac{3}{2} \sqrt{x} e^{x^{3/2}} \ln(x^3 + 1) + e^{x^{3/2}} \times \frac{3x^2}{x^3 + 1}$$

(b) Find $f'(1)$ for $f(x) = \frac{x-10}{(x^2+1)^2}$

$$f'(x) = \frac{(x^2+1)^2 \cdot 1 - (x-10) \cdot 2(x^2+1) \cdot 2x}{(x^2+1)^4}$$

Since $f'(1)$ is required, there is no need to simplify. Just substitute $x=1$.

$$f'(1) = \frac{(1+1)^2 - (1-10) \cdot 2(1+1) \cdot 2}{(1+1)^4} = \frac{4 + 72}{16} = \frac{19}{4}$$

(c) Use the chain rule to find $\frac{dy}{dx}$, if $y = \frac{u+1}{u-1}$, and $u = (x^2 + 5x - 1)^{1/4}$.

$$\frac{dy}{du} = \frac{(u-1) \cdot 1 - (u+1) \cdot 1}{(u-1)^2} = \frac{-2}{(u-1)^2}; \quad \frac{du}{dx} = \frac{1}{4} (x^2 + 5x - 1)^{-3/4} (2x + 5).$$

$$\frac{dy}{dx} = \frac{-2}{(u-1)^2} \times \frac{(2x+5)}{4(x^2+5x-1)^{3/4}} = \frac{-2(2x+5)}{4[(x^2+5x-1)^{1/4}-1]^2 (x^2+5x-1)^{3/4}}$$

(10%) 2. (a) Find $\int (\sqrt{\frac{7}{x}} + 2\sqrt{x} - 24x^2 - 3) dx = \int (\sqrt{7} x^{-1/2} + 2x^{1/2} - 24x^2 - 3) dx =$

$$\frac{\sqrt{7} x^{1/2}}{1/2} + \frac{2x^{3/2}}{3/2} - \frac{24x^3}{3} - 3x + C = 2\sqrt{7}x + \frac{4}{3}x^{3/2} - 8x^3 - 3x + C,$$

where C is a constant.

(b) If $g''(x) = 2 - \frac{4}{x^3}$, $g'(1) = 6$ and $g(1) = 13$, find $g(x)$.

We must integrate $g''(x)$ twice to find $g(x)$.

$$g''(x) = 2 - 4x^{-3} \Rightarrow g'(x) = \int (2 - 4x^{-3}) dx = 2x - \frac{4x^{-2}}{-2} + C_1$$

$$g'(x) = 2x + \frac{2}{x^2} + C_1 \text{ but } g'(1) = 6 \text{ so, } g'(1) = 2 + \frac{2}{1} + C_1 = 6$$

$$\Rightarrow C_1 = 2 + \left[g'(x) = 2x + \frac{2}{x^2} + 2 \right] \text{ Now } g(x) = \int (2x + 2x^{-2} + 2) dx =$$

$$x^2 + \frac{2x^{-1}}{-1} + 2x + C_2 \text{ but } g(x) = x^2 - \frac{2}{x} + 2x + C_2 \text{ but } g(1) = 13$$

$$\Rightarrow g(1) = 1 - 2 + 2 + C_2 = 13 \Rightarrow C_2 = 12$$

$$\boxed{g(x) = x^2 - \frac{2}{x} + 2x + 12}$$

(10%) 3. The frequency distribution for a random variable X is given below.

X	1	2	3	5
Frequency	20	40	80	60

(a) Construct the probability distribution of X .

Simply divide frequency of x by total sum of frequencies which is 200.


X	1	2	3	5
p(x)	$\frac{20}{200} = 0.1$	$\frac{40}{200} = 0.2$	0.4	0.3

(b) Find $p(X \leq 3) = p(X=3) + p(X=2) + p(X=1)$
 $= 0.4 + 0.2 + 0.1 = 0.7$


(c) Find $p(X > 5)$. X cannot be bigger than 5, so this event cannot happen \Rightarrow
 $p(X > 5) = 0$ (not ϕ : ϕ isn't a number just an empty set)

(15%) 4. The time it takes a college freshman to complete a certain reasoning test is normally distributed, with a mean of 30 minutes and standard deviation equal to 7 minutes.


(a) Find the probability that, a student chosen at random, will take between 15 and 25 minutes to finish the test.



(b) Find the probability that, a student, chosen at random, will take between 7 and 53 minutes to finish the test.




(c) Find the time t_1 below which only 30% of the students take to finish the test.



Area Under the Standard Normal Curve

	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.0000	.0040	.0080	.0120	.0160	.0199	.0239	.0279	.0319	.0359
0.1	.0398	.0438	.0478	.0517	.0557	.0596	.0636	.0675	.0714	.0753
0.2	.0793	.0832	.0871	.0910	.0948	.0987	.1026	.1064	.1103	.1141
0.3	.1179	.1217	.1255	.1293	.1331	.1368	.1406	.1443	.1480	.1517
0.4	.1554	.1591	.1628	.1664	.1700	.1736	.1772	.1808	.1844	.1879
0.5	.1915	.1950	.1985	.2019	.2054	.2088	.2123	.2157	.2190	.2224
0.6	.2257	.2291	.2324	.2357	.2389	.2422	.2454	.2486	.2518	.2549
0.7	.2580	.2612	.2642	.2673	.2704	.2734	.2764	.2794	.2823	.2852
0.8	.2881	.2910	.2939	.2967	.2995	.3023	.3051	.3078	.3106	.3133
0.9	.3159	.3186	.3212	.3238	.3264	.3289	.3315	.3340	.3365	.3389
1.0	.3413	.3438	.3461	.3485	.3508	.3531	.3554	.3577	.3599	.3621
1.1	.3643	.3665	.3686	.3708	.3729	.3749	.3770	.3790	.3810	.3830
1.2	.3849	.3869	.3888	.3907	.3925	.3944	.3962	.3980	.3997	.4015
1.3	.4032	.4049	.4066	.4082	.4099	.4115	.4131	.4147	.4162	.4177
1.4	.4192	.4207	.4222	.4236	.4251	.4265	.4279	.4292	.4306	.4319
1.5	.4332	.4345	.4357	.4370	.4382	.4394	.4406	.4418	.4429	.4441
1.6	.4452	.4463	.4474	.4484	.4495	.4505	.4515	.4525	.4535	.4545
1.7	.4554	.4564	.4573	.4582	.4591	.4599	.4608	.4616	.4625	.4633
1.8	.4641	.4649	.4656	.4664	.4671	.4678	.4686	.4693	.4699	.4706
1.9	.4713	.4719	.4726	.4732	.4738	.4744	.4750	.4756	.4761	.4767
2.0	.4772	.4778	.4783	.4788	.4793	.4798	.4803	.4808	.4812	.4817
2.1	.4821	.4826	.4830	.4834	.4838	.4842	.4846	.4850	.4854	.4857
2.2	.4861	.4864	.4868	.4871	.4875	.4878	.4881	.4884	.4887	.4890
2.3	.4893	.4896	.4898	.4901	.4904	.4906	.4909	.4911	.4913	.4916
2.4	.4918	.4920	.4922	.4925	.4927	.4929	.4931	.4932	.4934	.4936
2.5	.4938	.4940	.4941	.4943	.4945	.4946	.4948	.4949	.4951	.4952
2.6	.4953	.4955	.4956	.4957	.4959	.4960	.4961	.4962	.4963	.4964
2.7	.4965	.4966	.4967	.4968	.4969	.4970	.4971	.4972	.4973	.4974
2.8	.4974	.4975	.4976	.4977	.4977	.4978	.4979	.4979	.4980	.4981
2.9	.4981	.4982	.4982	.4983	.4984	.4984	.4985	.4985	.4986	.4986
3.0	.49865	.4987	.4987	.4988	.4988	.4989	.4989	.4989	.4990	.4990



(8%) 5. A box contains 5 red balls and 7 white balls. An experiment consists of drawing 6 balls at random from the box, with replacement. Find the probability that at least 4 of the balls are red.

This is a Bernoulli experiment where $n = 6$. But it success = getting red ball, then n . of successes $k \leq 5$ since we only have 5 red balls. $p = \frac{5}{12}$, $q = \frac{7}{12}$.

$$P(k \geq 4) = P(k=4) + P(k=5) = \binom{6}{4} \left(\frac{5}{12}\right)^4 \left(\frac{7}{12}\right)^2 + \binom{6}{5} \left(\frac{5}{12}\right)^5 \left(\frac{7}{12}\right)^1 = 0.197 \approx 0.20$$

(17%) 6. Of 140 students, 60 are studying French, 70 are studying Spanish, and 30 are studying both French and Spanish.

(a) A student is chosen at random. Find the probability that: $F =$ event student studying French.

(i) The student is studying neither French nor Spanish.

Sample Space




This is the complement of $F \cup S$.

$$P(F \cup S) = P(F) + P(S) - P(F \cap S) = \frac{60 + 70 - 30}{140} = \frac{100}{140}$$

$$1 - P(F \cup S) = 1 - \frac{100}{140} = \frac{40}{140} = \frac{2}{7} \quad (\text{these students are in shaded region})$$

(ii) The student is studying exactly one of the two languages.

Either French only or Spanish only (in shaded region).



$$P(F \text{ not } S) + P(S \text{ not } F) = \frac{30}{140} + \frac{40}{140} = \frac{70}{140} = \frac{1}{2}$$

(iii) The student is studying French, given that he is studying Spanish.

$$P(F|S) = \frac{P(F \cap S)}{P(S)} = \frac{3/140}{7/140} = \frac{3}{7}$$

(b) Let F be the event that the student is studying French, and S be the event that the student is studying Spanish. Are F and S statistically independent events? Give a clear reason for your answer.

F & S are statistically independent if & only if $P(F|S) = P(F)$.

$$P(F|S) = \frac{3}{7} \quad (\text{from (iii)}) \quad P(F) = \frac{60}{140} = \frac{3}{7} \Rightarrow$$

$P(F|S) = P(F)$ & the 2 events F & S are statistically independent.

(10%) 7. The scores on a quiz for a class of 50 students have a mean equal to 75 and a standard deviation equal to 9. If the teacher decides to raise each student's grade by 2 grades

(a) What will be the new mean (of the raised grades)? Justify your answer.

Let the old grade be x & the new grade be $x' = x + 2$.

$$\bar{x}' = \frac{\sum x'}{50} = \frac{\sum (x+2)}{50} = \frac{\sum x + 2 \times 50}{50}$$

$$= \frac{\sum x}{50} + 2 \Rightarrow \bar{x} + 2 = 75 + 2 = 77$$

The new mean is 77.

(b) What will be the new standard deviation? Justify your answer.

Let the new standard dev. be σ' .

$$\sigma' = \sqrt{\frac{\sum (x' - 77)^2}{50}}$$

but each x' increase by 2 so for each x' , the deviation from the mean $x' - 77 = x - 75$ since both x & mean increased by 2 $\Rightarrow \sigma' = 9$. No change in Standard Dev.

(18%) 8. A pair of dice, one red, one green, is rolled once. The random variable X is the maximum of the 2 numbers obtained on the faces of the dice.

(a) If A is the event that $X = 3$, list the set of outcomes belonging to this event.

$$X = 3 : \{(3, 3), (3, 2), (3, 1), (1, 3), (2, 3)\}$$

(b) Find the probability that $X \leq 3$.

$$X = 1 : \{(1, 1)\} \quad X = 2 : \{(1, 2), (2, 1), (2, 2)\}$$

$X = 3$: As in (a)

$$\text{so } P(X \leq 3) = \frac{\text{number of outcomes for } X \leq 3}{\text{total n. of outcomes (36)}}$$

$$P(X \leq 3) = \frac{1+3+5}{36} = \frac{9}{36} = \frac{1}{4}$$

(c) Find the probability that $X \neq 3$.

$$P(X=3) = \frac{5}{36} \quad P(X \neq 3) = 1 - \frac{5}{36} = \frac{31}{36}$$

(d) Let A be the event that $X = 3$, B be the event that X is even, and C be the event that $X \geq 4$.

(i) Are the events A, B and C mutually exclusive? Justify your answer.

2, 4, 6 belong to B.
4, 5, 6 " " C.
event
+ hat
 $B \cap C = \{X = 4, 6\}$ so $B \cap C \neq \emptyset$. B & C aren't mutually
exclusive \Rightarrow A, B, C are not either.

(ii) Are the events A, B and C collectively exhaustive? Justify your answer.

$A \cup B \cup C = \{X = 2, 3, 4, 5, 6\}$
 $X = 1 \notin A \cup B \cup C$. 1 is in sample
space, so A, B, C aren't collectively
exhaustive.