

---

---

**AMERICAN UNIVERSITY OF BEIRUT**  
**Faculty of Arts and Sciences**  
**Computer Science Department**

---

---

**CMPS 251**  
**FINAL EXAM**  
**FALL 2003-2004**  
Closed Book, Two hours

**GIVE YOUR ANSWERS ON THE QUESTION SHEET**  
**SUBMIT WITH BOOKLET**

<b>STUDENT NAME</b>	
<b>ID NUMBER</b>	

1. In this problem, we consider the following table for the function  $f(x)$ . All computations shall be carried out **with 8 significant figures**.

$x_i$	$y_i$
0.000	4.000000000000000
0.125	3.938461538461539
0.250	3.764705882352941
0.375	3.506849315068493
0.500	3.200000000000000
0.625	2.876404494382022
0.750	2.560000000000000
0.875	2.265486725663717
1.000	2.000000000000000

- (a) Write the polynomial of degree 3  $p(x)$  that would best approximate  $f(0.3)$ . Find  $p(0.3)$ .

- (b) Using the central difference formula to approximate  $f'(0.5)$ , followed by Richardson's extrapolation find the best approximation to  $f'(0.5)$ . For that purpose, fill out the following table.

$h$	$\phi_{c,h}$	$\phi_{c,h}^{(1)}$	$\phi_{c,h}^{(2)}$
0.5		×	×
		×	×
0.25			×
			×
0.125			

- (c) Using the Mid-point, trapezoidal and Simpson's rules followed by Romberg integrations fill out the following table used to approximate  $I = \int_0^1 f(x)dx$

$h$	$M_h$	$T_h$	$S_h$	$R_h^{(1)}$	$R_h^{(2)}$
$h_0$				×	×
				×	×
$\frac{h_0}{2}$					×
					×
$\frac{h_0}{4}$					
$\frac{h_0}{8}$	×		×	×	×
	×		×	×	×

2. Suppose a real number  $L$  is approximated by  $\phi(h)$  such that:

$$L = \phi(h) + c_1h^3 + c_2h^5 + c_3h^7 + \dots,$$

where the coefficients  $\{c_i\}$  are independent from  $h$ . What combination of  $\phi(h)$  and  $\phi(\frac{h}{2})$  would give a better approximation  $\phi^1(h)$  to  $L$  than  $\phi(h)$ ? What is the order  $\alpha$  of the approximation of  $L$  by  $\phi^1(h)$ , (i.e.  $L = \phi^1(h) + O(h^\alpha)$ )?

3. Loss of significant figures may result in the computation of the following functions of the variable  $x$  for certain values of  $x$ . Specify these values then propose alternative functions that would remedy the loss of significant figures. (If necessary you may use Taylor's series).

(a)  $f(x) = x + \sqrt{x^2 - 1}$

(b)  $g(x) = x - \sin(x)$

4. To perform Naive Gauss elimination for the following **quadridiagonal matrix**

$$\begin{pmatrix} d_1 & u_1 & v_1 & 0 & .. & .. & 0 & 0 \\ l_1 & d_2 & u_2 & v_2 & 0 & .. & .. & 0 \\ 0 & l_2 & d_3 & u_3 & v_3 & 0 & .. & 0 \\ 0 & 0 & l_3 & d_4 & u_4 & v_4 & .. & 0 \\ \vdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ 0 & .. & 0 & l_i & d_i & \cdot & \cdot & \cdot \\ \vdots & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \vdots & \cdot & \cdot & \cdot & l_{n-3} & d_{n-2} & u_{n-2} & v_{n-2} \\ 0 & \cdot & \cdot & \cdot & \cdot & l_{n-2} & d_{n-1} & u_{n-1} \\ 0 & .. & .. & 0 & .. & 0 & l_{n-1} & d_n \end{pmatrix}$$

One uses the following algorithm:

for i from 1 to n-1

$$l_i = l_i/d_i$$

$$d_{i+1} = d_{i+1} - l_i * u_i$$

if i < n-1

$$u_{i+1} = u_{i+1} - l_i * v_i$$

end

end

Give the exact number of **floating point operations** needed to perform this algorithm.

5. Give with justification the minimum number of arithmetic operations (additions, subtractions and multiplications) to compute, using nested multiplication, the following polynomials

Polynomial $p(x)$	Minimum number of arithmetic operations
$(x - 2)^{17} + (x - 2)^{31}$	
$4x^5 - 6x^{12} + 2x^{17} - x^{33}$	