

3.12 This problem asks that we calculate the theoretical densities of Al, Ni, Mg, and W.

Since Al has an FCC crystal structure, $n = 4$, and $V_C = 16R^3\sqrt{2}$ (Equation 3.4). Also, $R = 0.143 \text{ nm}$ ($1.43 \times 10^{-8} \text{ cm}$) and $A_{\text{Al}} = 26.98 \text{ g/mol}$. Employment of Equation 3.5 yields

$$\begin{aligned}\rho &= \frac{nA_{\text{Al}}}{V_C N_A} \\ &= \frac{(4 \text{ atoms/unit cell})(26.98 \text{ g/mol})}{\left\{ \left[(2)(1.43 \times 10^{-8} \text{ cm})(\sqrt{2}) \right]^3 / (\text{unit cell}) \right\} (6.023 \times 10^{23} \text{ atoms/mol})} \\ &= 2.71 \text{ g/cm}^3\end{aligned}$$

The value given in the table inside the front cover is 2.71 g/cm^3 .

Nickel also has an FCC crystal structure and therefore

$$\begin{aligned}\rho &= \frac{(4 \text{ atoms/unit cell})(58.69 \text{ g/mol})}{\left\{ \left[(2)(1.25 \times 10^{-8} \text{ cm})(\sqrt{2}) \right]^3 / (\text{unit cell}) \right\} (6.023 \times 10^{23} \text{ atoms/mol})} \\ &= 8.82 \text{ g/cm}^3\end{aligned}$$

The value given in the table is 8.90 g/cm^3 .

Magnesium has an HCP crystal structure, and from the solution to Problem 3.6,

$$V_C = \frac{3\sqrt{3} a^2 c}{2}$$

and, since $c = 1.624a$ and $a = 2R = 2(1.60 \times 10^{-8} \text{ cm}) = 3.20 \times 10^{-8} \text{ cm}$

$$V_C = \frac{(3\sqrt{3})(1.624)(3.20 \times 10^{-8} \text{ cm})^3}{2} = 1.38 \times 10^{-22} \text{ cm}^3/\text{unit cell}$$

Also, there are 6 atoms/unit cell for HCP. Therefore the theoretical density is

$$\rho = \frac{nA_{\text{Mg}}}{V_C N_A}$$