

N.D.U

MAT 112
First Test

Name: _____

ID #: _____

Section: A

Time: 60 minutes

Directions

1. Write neatly and clearly.
2. Do not use pencils unless for graphing.
3. Show all work.
4. No calculators are allowed.
5. Turn off your mobile phones.

96

I. Evaluate $\int 4 \sec\left(\frac{x}{2}\right) \cdot \tan\left(\frac{x}{2}\right) dx$. (10 points)

$$= 4 \int \sec\left(\frac{x}{2}\right) \cdot \tan\left(\frac{x}{2}\right) dx = 4 \times \frac{1}{\frac{1}{2}} \left(\sec \frac{x}{2}\right) = \boxed{8 \sec \frac{x}{2} + C}$$

II. Evaluate $\int x^{1/3} \sin(x^{4/3} - 8) dx$. (8 points)

Let $u = x^{4/3} - 8$ $du = \frac{4}{3} x^{4/3 - 1} dx = \frac{4}{3} x^{1/3} dx \Rightarrow \frac{3}{4} du = x^{1/3} dx$

So $\int x^{1/3} \sin(x^{4/3} - 8) dx = \frac{3}{4} \int du \sin u = \boxed{\frac{3}{4} (-\cos)(x^{4/3} - 8) + C}$

23
9
11
12

III. Solve the following initial value problem:

$$\frac{d^2y}{dx^2} = 2 - 6x; \quad y'(0) = 4 \text{ and } y(0) = 1$$

(12 points)

$$\int \frac{d^2y}{dx^2} dx = \int (2 - 6x) dx = \frac{dy}{dx} = 2x - \frac{6x^2}{2} + C = 2x - 3x^2 + C$$

① $y'(0) = 4$ so $4 = 0 - 0 + C$ $C = 4$ then $\frac{dy}{dx} = 2x - 3x^2 + 4$

$$\int \frac{dy}{dx} \cdot dx = \int (2x - 3x^2 + 4) dx = y = \frac{2x^2}{2} - \frac{3x^3}{3} + 4x + C = x^2 - x^3 + 4x + C$$

$$= x^2 - x^3 + 4x + C$$

$y(0) = 1$ so $1 = 0 - 0 + 0 + C$ $C = 1$ ✓



then $y = x^2 - x^3 + 4x + 1$ ✓

IV. Graph the integrand and use area to evaluate the integral: $\int_{-4}^0 \sqrt{16-x^2} dx$

$$y = \sqrt{16-x^2} \quad y^2 = 16-x^2 \Rightarrow x^2 + y^2 = 16 \quad (10 \text{ points})$$

It is the equation of the circle with center is $(0,0)$ and radius 4.

The required area is between -4 and 0 $[-4,0]$ and we have $y = \sqrt{16-x^2}$ then:



$$\int_{-4}^0 \sqrt{16-x^2} dx = \frac{1}{4} \pi R^2 = \frac{1}{4} \pi 16 = 4\pi \text{ u}^2 = \text{Area of } \frac{1}{4} \text{ of the circle}$$

VII. Given $y = \int_0^{x^2} \cos(\sqrt{t}) dt$. Find $\frac{dy}{dx}$.

(10 points)

$$\frac{dy}{dx} = \frac{d}{dx} \int_0^{x^2} \cos \sqrt{t} dt$$

let $u = x^2 \quad du = 2x dx$

$$= \frac{d}{dx} \cdot \frac{du}{dx} = 2x \cdot \cos \sqrt{x^2} = \boxed{2x \cos |x| = \frac{dy}{dx}} =$$

$$= 2x \cos |x|,$$

since $\cos(-x) = \cos(x)$,

VIII. Find the area of the following shaded region:

(14 points)

(14) In A: ~~Upper~~ curve = $f(x) = 2x^3 - x^2 - 5x = y_1$
 Lower " = $g(x) = -x^2 + 3x = y_2$


$$f(x) - g(x) = 2x^3 - x^2 - 5x + x^2 - 3x = 2x^3 - 8x$$

$$\text{Area of A} = \int_{-2}^0 f(x) - g(x) dx$$

$$= \int_{-2}^0 2(2x^3 - 8x) dx$$

$$= \left[\frac{2x^4}{4} - \frac{8x^2}{2} \right]_{-2}^0 = \left[\frac{x^4}{2} - 4x^2 \right]_{-2}^0$$

$$0 - \left(\frac{16}{2} - 16 \right) = -(8 - 16) = 8$$



In B, Lower curve $2x^3 - x^2 - 5x = g(x) = y_1$
 Upper curve $-x^2 + 3x = f(x) = y_2$

$$f(x) - g(x) = -x^2 + 3x - 2x^3 + x^2 + 5x = -2x^3 + 8x$$

$$\text{Area of B} = \int_0^2 (-2x^3 + 8x) dx = \left[-\frac{2x^4}{4} + \frac{8x^2}{2} \right]_0^2 = \left[-\frac{x^4}{2} + 4x^2 \right]_0^2$$

$$= -\frac{16}{2} + 16 - 0 = 8$$

$$\text{Total Area} = \text{Area of A} + \text{Area of B} = 8 + 8 = \boxed{16 \text{ m}^2}$$

IX. Consider a solid that lies between planes perpendicular to the x -axis at $x = 0$ and $x = 4$. The cross sections of the solid are perpendicular to the x -axis and are between these planes, and run from the parabola $y = -\sqrt{x}$ to the parabola $y = \sqrt{x}$. These cross sections are circular disks with diameters in the xy -plane. Find a formula for the area $A(x)$ of the cross sections, then use it to get the volume V of the solid.

(18)

$$AB = 2R = 2\sqrt{x} = \text{diameter} \quad \checkmark$$

$$R = \frac{2\sqrt{x}}{2} = \sqrt{x} \quad \checkmark$$

$$A(x) = \text{Area of the circle} \quad \checkmark$$

$$= \pi R^2 = \pi x \quad \checkmark$$

Volume of the solid

$$V = \int_0^4 A(x) dx = \int_0^4 (\pi x) dx \quad \checkmark$$

$$= \pi \int_0^4 x dx = \pi \left[\frac{x^2}{2} \right]_0^4 \quad \checkmark$$

$$= \pi \left(\frac{16}{2} \right) = \boxed{8\pi} \quad \checkmark$$

