

Name: Signature:

Student number:

Section number (Encircle): 3      5      6

**1. Instructions:**

- No calculators are allowed.
- Make sure that you have exactly 20 questions.
- Each question has only one correct answer.
- Make sure to circle your answers appropriately below.

**2. Grading policy:**

- 5 points for each correct answer.
- -1 point (penalty) for each wrong answer.
- 0 point for zero or more than one answer.

**Circle the appropriate answers:**

1. a	b	c	d	11. a	b	c	d
2. a	b	c	d	12. a	b	c	d
3. a	b	c	d	13. a	b	c	d
4. a	b	c	d	14. a	b	c	d
5. a	b	c	d	15. a	b	c	d
6. a	b	c	d	16. a	b	c	d
7. a	b	c	d	17. a	b	c	d
8. a	b	c	d	18. a	b	c	d
9. a	b	c	d	19. a	b	c	d
10. a	b	c	d	20. a	b	c	d

(0)

1. The area of the region lying inside the circle  $r = -3 \cos \theta$  and outside  
[www.amal-aub.org](http://www.amal-aub.org)

the cardioid  $r = 1 - \cos \theta$  is

(a)  $3(\pi + 3\sqrt{2} - 1)$ .

(b)  $\pi$ .

(c)  $2\pi$ .

(d)  $2(\pi + 2\sqrt{3} - 2)$ .

2. The points of intersection of the curves  $r = 2$  and  $r = 4 \cos 2\theta$  are

(a)  $(2, 0), (2, \pi), (2, \pm\pi/2)$ .

(b)  $(2, \pm\pi/6), \pm\pi/3, (2, \pm 2\pi/3), (2, \pm 5\pi/6)$ .

(c)  $(2, \pm\pi/12), \pm 5\pi/12, (2, \pm 13\pi/12), (2, \pm 17\pi/12)$ .

(d)  $(2, \pm\pi/4), (2, \pm 5\pi/4)$ .

3. The surface  $x^2/16 - y^2/25 = z/4$  is

(a) a circular paraboloid.

(b) a hyperbolic paraboloid.

(c) an elliptic cone.

(d) a one-sheeted hyperboloid.

4. If  $f(x, y) = x^2y^3/[x^8 + y^4]$  for  $(x, y) \neq (0, 0)$  and  $f(0, 0) = 0$ , then

(a)  $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = 1/2$ .

(b)  $f$  is discontinuous at  $(0,0)$ .

(c)  $\lim_{(x,y) \rightarrow (0,0)} f(x, y) = 0$ .

(d)  $f$  is continuous at  $(0,0)$ .

(1)

5. An estimate of the error obtained by taking the first four terms of the  
[www.amal-aub.org](http://www.amal-aub.org)

series

$$\sum_{n=1}^{\infty} (-1)^{n+1} (0.01)^n / n^3 \quad \text{is}$$

(a)  $1.56 \times 10^{-10}$ .

(b)  $8 \times 10^{-11}$ .

(c)  $1.56 \times 10^{-12}$ .

(d)  $8 \times 10^{-13}$ .

6. The interval of convergence of the power series

$$\sum_{n=0}^{\infty} 2^n (x - 3)^n / n^2 \quad \text{is}$$

(a)  $5/2 < x < 7/2$ .

(b)  $1 \leq x \leq 5$ .

(c)  $5/2 \leq x \leq 7/2$ .

(d)  $1 < x < 5$ .

7. The Maclaurian series of the integral

$$\int_0^x \frac{\sin t^4}{t^2} dt \quad \text{is}$$

(a)  $\sum_{n=0}^{\infty} (-1)^n x^{8n+1} / (8n+1)[(2n+1)!]$ .

(b)  $\sum_{n=0}^{\infty} (-1)^n x^{8n+3} / (8n+3)[(2n+1)!]$ .

(c)  $\sum_{n=0}^{\infty} (-1)^n x^{8n+1} / (8n+3)[(2n+1)!]$ .

(d)  $\sum_{n=0}^{\infty} (-1)^n x^{8n+3} / (8n+3)[(2n+3)!]$ .

(2)

8. If  $a_n = (\frac{3}{5})^n + \frac{4^n}{n!}$  and  $b_n = n(e^{\frac{1}{n}} - 1)$ , then  
[www.amal-aub.org](http://www.amal-aub.org)

- (a) the sequences  $\{a_n\}$  and  $\{b_n\}$  diverge.
- (b) the sequences  $\{a_n\}$  and  $\{b_n\}$  converge.
- (c) the sequence  $\{a_n\}$  diverges and  $\{b_n\}$  converges.
- (d) the sequence  $\{a_n\}$  converges and  $\{b_n\}$  diverges.

9. If  $a_n = \frac{2n+1}{n^2(n+1)^2}$  and  $b_n = \frac{1}{n(\ln n)^{ln 2}}$ , then

- (a) the series  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  diverge.
- (b) the series  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  converge.
- (c) the series  $\sum_{n=1}^{\infty} a_n$  diverges and  $\sum_{n=1}^{\infty} b_n$  converges.
- (d) the series  $\sum_{n=1}^{\infty} a_n$  converges and  $\sum_{n=1}^{\infty} b_n$  diverges.

10. If  $a_n = (1 - \frac{1}{n})^n$  and  $b_n = \frac{\cos n\pi}{\sqrt{n}}$ , then

- (a) the series  $\sum_{n=1}^{\infty} a_n$  diverges and  $\sum_{n=1}^{\infty} b_n$  converges.
- (b) the series  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  converge.
- (c) the series  $\sum_{n=1}^{\infty} a_n$  and  $\sum_{n=1}^{\infty} b_n$  diverge.
- (d) the series  $\sum_{n=1}^{\infty} a_n$  converges and  $\sum_{n=1}^{\infty} b_n$  diverges.

11. The Taylor series of  $\ln x$  at  $x = 3$  is

- (a)  $\ln 3 + \sum_{n=1}^{\infty} (-1)^n(x - 3)^n/n3^n$ .
- (b)  $\ln 3 + \sum_{n=1}^{\infty} (-1)^{n+1}(x - 3)^n/n!3^n$ .
- (c)  $\ln 3 + \sum_{n=1}^{\infty} (-1)^{n+1}(x - 3)^n/n3^n$ .
- (d)  $\ln 3 + \sum_{n=1}^{\infty} (-1)^n(x - 3)^n/n!3^n$ .

(3)

12. If  $z = f(x, y)$  where  $x = r \cos \theta$  and  $y = r \sin \theta$ , then  
[www.amal-aub.org](http://www.amal-aub.org)

- (a)  $f_x^2 + f_y^2 = r^2 f_r^2 + f_\theta^2$ .
- (b)  $f_x^2 + f_y^2 = f_r^2 + r^{-2} f_\theta^2$ .
- (c)  $f_x^2 + f_y^2 = f_r^2 + f_\theta^2$ .
- (d)  $f_x^2 + f_y^2 = f_r^2 - f_\theta^2$ .

13. An equation of the tangent plane to the surface  $x^2 + 2xy + 2y^2 + z = 12$

at the point  $P(2, -3, 2)$  is

- (a)  $-8x + 18y - z = -72$ .
- (b)  $2x + 8y - z = -22$ .
- (c)  $2x + 18y - z = -52$ .
- (d)  $-8x + 8y - z = -42$ .

14. The tangent line to the curve of intersection of the paraboloid  $x^2 + y^2 - z = 0$  and the ellipsoid  $2x^2 + 3y^2 + z^2 - 9 = 0$  through the point  $P(1, -1, 2)$  has parametric equations:

- (a)  $x = 1 + 7t, y = -1 + 6t, z = 2 + 2t$ .
- (b)  $x = 1 + 5t, y = -1 - 7t, z = 2 + t$ .
- (c)  $x = 1 - 3t, y = -1 + 9t, z = 2 + 3t$ .
- (d)  $x = 1 + 4t, y = -1 - 7t, z = 2 - t$ .

(4)

15. The rate of change of the temperature function  $T(x, y) = 4x^2 - y^2 + 16z^2$  at the point  $P(-1, 2, 1)$  is zero in the direction of the vector:

(a)  $\langle 3, 4, -2 \rangle$ .

(b)  $\langle 1, -1, 2 \rangle$ .

(c)  $\langle 1, -2, 0 \rangle$ .

(d)  $\langle 0, 8, -1 \rangle$ .

16. The function  $f(x, y) = 3x^2 - 4xy + 5y^2 - 7$  admits -7 as a

(a) local maximum value.

(b) an absolute minimum value.

(c) a value at a saddle point.

(d) an absolute maximum value.

17. The value of the double integral

$$\int_0^2 \int_{y/2}^1 ye^{x^3} dx dy \quad \text{is}$$

(a)  $(e - 1)/4$ .

(b)  $5(e - 1)/4$ .

(c)  $2(e - 1)/3$ .

(d)  $(e - 1)/3$ .

(5)

18. The average value of the function  $e^{-(x^2+y^2)}$  over the first quadrant of  
[www.amal-aub.org](http://www.amal-aub.org)

the disc  $x^2 + y^2 \leq 1$  is

(a)  $(e - 1)/2$ .

(b)  $e - 1$ .

(c)  $1 - e^{-1}$ .

(d)  $(1 - e^{-1})/2$ .

19. The volume of the solid bounded by the paraboloids  $z = 2x^2 + y^2$

and  $z = 12 - x^2 - 2y^2$  is

(a)  $20\pi$ .

(b)  $24\pi$ .

(c)  $18\pi$ .

(d)  $22\pi$ .

20. The solid whose volume is given by the triple integral in spherical coordinates as

$$\int_0^{2\pi} \int_0^{\pi/6} \int_0^{2\cos\phi} \rho^2 \sin\phi d\rho d\phi d\theta$$

is bounded by

(a) a cone and a cylinder.

(b) a sphere and a cylinder.

(c) a cone and a sphere.

(d) a sphere and a paraboloid.

(6)